

2026 CLS Quicksheet

CLS Fungicide Program BMPs

- **Start early when the leaves between rows are ~4 inches apart, GDU's reach 1500 from your planting date, or by June 21st.** An early application provides significantly better control than fungicide programs that start later. Delaying the onset of disease is critical and early season testing shows CLS infections ramping up by the 3rd week of June.
- Use a **spray volume of 20 to 25 gpa** (5 to 7 gpa for aerial applications).
- Utilize nozzle selection, pressure, and deposition aid adjuvants to achieve **optimal coverage**.
 - A medium droplet size will provide the best coverage in the most situations (250-300 microns).
- If possible, avoid making applications during periods of high temperature or wind speeds over 10mph. These conditions will negatively impact coverage and penetration into the canopy.
- Reducing ground speed and maintaining proper boom height can also improve coverage.
- If a scheduled fungicide application coincides with a rain event, make the fungicide application ahead of the rain. This will kill the CLS that is present that would otherwise sporulate after the rain event prior to the field being fit to spray again.
- **Make fungicide applications on a 10-day interval for all varieties, unless environmental conditions dictate otherwise.**
 - CLS DIV's will begin posting in June.
<https://www.smbsc.com/agronomy/AgronomyDefault>
- Maintaining protection into September is important as the new DIV matrix being developed indicates that free water is important for CLS infection and September has the potential for a high occurrence of overnight dew.
- SDHI fungicide products are not considered effective tank mix partners. However, early season leaf sample testing indicates that the CLS population is susceptible to benzimidazoles (Topsin) and Qols (Headline, Priaxor, etc) at least during the early part of the growing season.
- There is cross resistance between Inspire XT and Provysol, and also between Proline/Phobos, Minerva/Domark, Enable, and Topguard. Products from these two groups should be rotated during the season.
- Do not tank mix fungicides and glyphosate.
- Consult labels before tank-mixing fungicides with other crop protection products.
- Always follow the label for application instructions and restrictions.
- **Please follow the CLS Program outlined below.**

CLS Fungicide Program

Start when leaves are ~4" apart or 1500 GDU
Applications on 10-day intervals

1. Proline* + EBDC
2. Tin + Topsin*
3. Qol (strobi) + EBDC
4. Tin + EBDC
5. Inspire XT** + EBDC
6. Tin + EBDC
7. DMI Group 1*** + EBDC



*Or generic

**DMI Group 2: Inspire XT and Provysol. Or generic. Regev and Luna Flex have a 7-day PHI compared to Inspire XT 21-day PHI.

***DMI Group 1: Domark/Minerva, Lucento/Topguard, Enable, and Proline.

CLS Fungicide Product Options

Page 2

DMI Products	Rate/Acre	REI	PHI
Domark	6.9 oz	12hrs	14 days
Minerva	13 oz	12hrs	14 days
Proline 480 SC or Prozio SC	5.7 oz	12hrs	7 days
Enable 2F	8 oz	12hrs	14 days
Phobos FC	7.6 oz	12hrs	7 days
Topguard	14 oz	12hrs	21 days
Provysol	5 oz	12hrs	7 days
InspireXT or EsquireXT	7 oz	12hrs	21 days
*Lucento Topguard+SDHI	5.5 oz	12hrs	21 days
*Luna Flex Inspire+SDHI	13.6 oz	12hrs	7 days
*REGEV Inspire+TTO	8.5 oz	12hrs	7 days

*Cross resistance has been documented between Proline/Phobos, Minerva/Domark, Enable, and Topguard products (DMI Group 1, gold font), and between Inspire XT and Provysol (DMI Group 2, black font). Rotate the use of these two groups in season. We should only use each product once during a season.

*These premixed fungicide products do not contain two effective modes of action. If these products are used, they should be tank mixed with an EBDC.

*Be aware of which triazole is in the premix so that we are not applying more than one triazole from a resistance group per season.

*Other premixes are available for use in sugar beets. However, these products do not align with the program goals.

QoI Products	Rate	REI	PHI
Gem 500 SC / Flint Extra	3.6 oz	12 hrs	21 days
Headline	9 oz	12 hrs	7 days
Priaxor Headline+SDHI	6.7 oz	12 hrs	7 days

*There may be other QoI products available for use in sugar beets. Follow label instructions.

The information contained in this quicksheet is meant to provide options for a CLS fungicide program for your operation in 2026. However, it can not provide all the details for every application. Consult your agriculturalist and the product label.

Copper	Rate	REI	PHI
Badge SC	2 pts	48hrs	0 day
Champ 2 Flowable	2 pts		
Cuprofix Flex	2 lbs		
Kocide 3000	1.25 lbs		

*There may be other copper products available for use in sugar beets. Follow label instructions.

*Do not apply glyphosate within 7 to 10 days of a copper product as this may cause phytotoxicity.

Benzimidazole	Rate	REI	PHI
T-Methyl 4.5 F	10 oz	24hrs	21 days
Topsin 4.5 FL			
Incognito 4.5 F			
Talaris 4.5 F			
Topsin M WSB	0.5 lb		
T-Methyl 70 WSB			
Talaris 70 WSB			
Incognito 85 WDG			

*There may be other products available for use in sugar beets. Follow label instructions.

Tin Products	Rate	REI	PHI
Super Tin 4L	8 oz	48hrs	7 days
Agri Tin Flowable	8 oz		

*Do not exceed 24 oz/A of Super Tin 4L/Agri Tin Flowable per season.

EBDC Products	Rate	REI	PHI
Manzate Max / Dithane F-45	1.6 qts	24hrs	14 days
Manzate Pro-Stick / Koverall / Dithane M45 / Penncozeb 80WP or 75DF	2 lbs		

*There may be other EBDC products available for use in sugar beets. Follow label instructions.

**Agricultural Department
Southern Minnesota
Beet Sugar Cooperative**

Updated Feb 4th, 2026



Cercospora Leaf Spot Rating Scale

